

Acknowledgement

We would like to thank Clinical Research Development Center of Ghaem Hospital for their assistant in this manuscript. This study was supported by a grant from the Vice Chancellor for Research of the Mashhad University of Medical Sciences for the research project as a medical student thesis with approval number of 910349.

Conflict of Interest

The authors declare no conflict of interest.

References

1. Berg AT, Shinnar S. The risk of seizure recurrence following a first unprovoked seizure: a quantitative review. *Neurology*. 1991;41:965-972.
2. Olafsson E, Ludvigsson P, Gudmundsson G, et al. Incidence of unprovoked seizures and epilepsy in Iceland and assessment of the epilepsy syndrome classification: a prospective study. *Lancet Neurol*. 2005;4:627-634.
3. Hermann B, Seidenberg M. Neuropsychology and temporal lobe epilepsy. *CNS Spectr*. 2002;7:343-348.
4. Kramer U, Rivello JJ, Carmat P. Clinical characteristics of complex partial seizures: a temporal versus a frontal lobe onset. *Seizure*. 1997;6:57-61.
5. Oyegbile TO, Bhattacharya A, Seidenberg M, et al. Quantitative MRI biomarkers of cognitive morbidity in temporal lobe epilepsy. *Epilepsia*. 2006;47:143-152.
6. Seidenberg M, Kelly KG, Parrish J, et al. Ipsilateral and contralateral MRI volumetric abnormalities in chronic unilateral temporal lobe epilepsy and their clinical correlates. *Epilepsia*. 2005;46:420-430.
7. Milesi G, Garbelli R, Zucca I, et al. Assessment of human hippocampal developmental neuroanatomy by means of ex-vivo 7T magnetic resonance imaging. *Int J Dev Neurosci*. 2014;34C:33-41.
8. Bote RP, Blázquez-Llorca L, Fernández-Gil MA, et al. Hippocampal sclerosis: histopathology substrate and magnetic resonance imaging. *Semin Ultrasound CT MR*. 2008;29:2-14.
9. Scott RC, King MD, Gadian DG, et al. Hippocampal abnormalities after prolonged febrile convulsion: a longitudinal MRI study. *Brain*. 2003;126:2551-2557.
10. Lawson J, Vögrin, S, Bleasel A, et al. Cerebral and cerebellar volume reduction in children with intractable epilepsy. *Epilepsia*. 2000;41:1456-1462.
11. Berg AT, Pardoe HR, Fulbright RK, et al. Hippocampal size anomalies in a community-based cohort with childhood-onset epilepsy. *Neurology*. 2011;76:1415-1421.
12. Luby M, Spencer DD, Kim JH, et al. Hippocampal MRI volumetrics and temporal lobe substrates in medial temporal lobe epilepsy. *Magn Reson Imaging*. 1995;13:1065-1071.
13. Daley M, Ott D, Blanton R, et al. Hippocampal volume in childhood complex partial seizures. *Epilepsy Res*. 2006;72:57-66.
14. Provenzale JM, Barboriak DP, VanLandingham K, et al. Hippocampal MRI signal hyperintensity after febrile status epilepticus is predictive of subsequent mesial temporal sclerosis. *Am J Roentgenol*. 2008;190:976-983.
15. Cao QJ, Wang H. Changes in the hippocampal volume in the acute phase following febrile convulsions in children. *Zhongguo Dang Dai Er Ke Za Zhi*. 2010;12:259-261.
16. Yoong M, Martinos MM, Chin RF, et al. Hippocampal volume loss following childhood convulsive status epilepticus is not limited to prolonged febrile seizures. *Epilepsia*. 2013;54:2108-2115.
17. Moore KR, Swallow CE, Tsuruda JS. Incidental detection of hippocampal sclerosis on MR images: is it significant? *Am J Neuroradiol*. 1999;20:1609-1612.