



Original research article

Evaluation of the serum prooxidant-antioxidant balance before and after vitamin D supplementation in adolescent Iranian girls

Ameneh Timar^{a,1}, Maryam Saberi-Karimian^{b,1}, Hamideh Ghazizadeh^b,
Seyed Mohammad Reza Parizadeh^b, Reihaneh Sabbaghzadeh^c, Maryam Emadzadeh^d,
Fatemeh Eshaghi^e, Shima Tavallaie^b, Gordon A. Ferns^f, Majid Ghayour-Mobarhan^{b,g,*}

^a Department of Modern Sciences and Technologies, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran

^b Metabolic Syndrome Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

^c Department of Biology, Hakim Sabzevari University, Sabzevar, Iran

^d Clinical Research Unit, Mashhad University of Medical Sciences, Mashhad, Iran

^e Faculty of Basic Sciences, Hakim Sabzevari University, Sabzevar, Iran

^f Brighton & Sussex Medical School, Division of Medical Education, Falmer, Brighton, UK

^g Cardiovascular Research Center, Mashhad University of Medical Sciences, Mashhad, Iran

ARTICLE INFO

Keywords:

Prooxidant- antioxidant balance

ROS

25OHD

Vitamin D deficiency

ABSTRACT

Purpose: Oxidative stress is caused by an imbalance between the antioxidant defenses and pro-oxidant production in favor of pro-oxidant production. Vitamin D has the potential for both pro- and anti-oxidative effects. The aim of this study was to assess the effect of high dose vitamin D supplementation on the prooxidant-antioxidant balance (PAB) in Iranian girls attending High School.

Materials and methods: A total of 464 girls aged 12–18 years were asked to take vitamin D capsules containing 50000IU vitamin D₃ once a week for a period of 9 weeks. All variables were determined at baseline and after 9 weeks of intervention. Fasting blood samples were taken from all subjects. The serum levels of 25OHD were measured using an electrochemiluminescence method. Serum PAB levels were determined using an ELISA reader at a wavelength of 450 nm.

Results: Vitamin D supplementation was associated with an increase in serum PAB ($P < 0.001$) and a reduction in serum LDL-C ($P < 0.001$), total cholesterol ($P < 0.001$) and HDL-C ($P < 0.01$) serum levels in Iranian adolescent girls. The results obtained from the current study show that there were significant improvements in weight ($P < 0.001$), BMI ($P < 0.001$) and FBG ($P = 0.02$) in adolescent girls who had 50–74.9 nmol/L serum 25OHD levels compared to < 50 nmol/L ones after the vitamin D supplementation. There was no significant association between the serum PAB and all biochemical factors ($P > 0.05$ for all variables).

Conclusions: The results showed that vitamin D supplementation has increased the PAB levels in teenage girls.

1. Introduction

Vitamin D (25-hydroxyvitamin D or 25OHD) is a fat-soluble steroid hormone which is often ingested from dietary sources (fatty fish and dairy products), but is also synthesized in the skin following exposure to UVB light [1]. In addition to its well known role in bone health, 25OHD can also protect against the occurrence of many types of cancers in humans [2–4]. 25OHD is involved in calcium homeostasis and takes part in several cellular processes, for example metabolism, proliferation, and differentiation, by both non-genomic function and its known

receptors (VDR)-mediated transcriptional effects [5].

About one third to half of the adult population globally (both in developed and developing countries) are affected by vitamin D deficiency [6,7]; in Canada and the USA more than two-thirds of individuals have a suboptimal serum vitamin D concentrations [8,9]. Reports indicate that about 79–81 % of adolescents in Iran are vitamin D deficient [10,11]. Studies in Mashhad have shown a prevalence of 80% deficiency and 12% insufficiency in vitamin D levels [12].

Oxidative stress occurs when there is an imbalance between the production of antioxidant defenses and pro-oxidant production in favor

* Corresponding author at: Metabolic Syndrome Research Center, Faculty of Medicine, Mashhad University of Medical Science, Vakil Abad Blvd., Opposite to Mellat Park, Mashhad 99199-91766, Iran. Tel.: +985138002288; fax: +985138002287.

E-mail address: ghayourm@mums.ac.ir (M. Ghayour-Mobarhan).

¹ Equal first author.

<https://doi.org/10.1016/j.advms.2018.10.004>

Received 21 February 2018; Accepted 31 October 2018

1896-1126/ © 2018 Medical University of Białystok. Published by Elsevier B.V. All rights reserved.

of pro-oxidant production [13,14]. There is a fine balance between the elimination and the production of the reactive oxygen species (ROS) [15]. Some studies have shown that the serum prooxidant-antioxidant balance (PAB) assay values are increased in patients with coronary artery disease and type II diabetes, and that the PAB assay can provide a simple surrogate measure of ROS [15–17]. The cellular oxidative condition is controlled by adjusting the level of antioxidant amount/activity and the level of oxidative source exposure/activity [18]. Therefore, reducing exposure to xenobiotics and radiation, suppressing the inflammatory response, suppressing enzyme activity and uncoupling mitochondrial potential can reduce oxidative stress [18].

The antioxidant properties of 25OHD were first reported in an *in vitro* model of ROS-induced neurotoxicity, and it was found to act by regulating the proteins that can reduce oxidative stress [19]. 25OHD can increase glutathione and protect neurons from oxidative degenerative processes [20]. 25OHD can also inhibit iron-related lipid peroxidation of membranes which may prevent free radical-induced cell membrane damage [21]. However, the mechanism of the vitamin D effects on the oxidative stress are not entirely clear. There is some evidence that the effects of 25OHD in breast cancer cells are caused by reducing antioxidant capacity and increasing ROS, and that it can potentiate the effects of ROS-producing drugs [22–25]. The aim of this study was to investigate the effects of a 3-month intervention with high dose vitamin D supplementation on a measure of the balance between serum pro- and anti-oxidants, measured using the PAB assay.

2. Materials and methods

2.1. Study population and design

The study population comprises adolescent girls who were recruited as a part of a previous study as described by Khayatzadeh et al. [26], from the cities of Sabzevar and Mashhad, in northeastern Iran from among girls aged 12–18 years. Informed consent was acquired from all participants. The study protocol is approved by ethical committee of Mashhad University of Medical Science (MUMS), Iran (IR-MUMS.fm.REC.1395.12). Inclusion and exclusion criteria were described previously [26]. A total of 464 girls aged 12–18 years were asked to take vitamin D capsules containing 50000IU vitamin D₃ once a week for a period of 9 weeks.

2.2. Measurements

2.2.1. Clinical and biochemical features

Anthropometric indices were determined at baseline and after 9 weeks of intervention using standard procedures. Systolic (SBP) and diastolic (DBP) blood pressures were measured using a standard mercury sphygmomanometer on the left arm in the sitting position following a 15 min rest by a standard procedure. Fasting blood samples were taken after 14 h overnight fast from each subject and collected into vacuum tubes in the morning. The serum of blood samples was separated using centrifugation (Hettich model D-78532, New York, USA) and aliquots of serum were frozen at -80°C for future analysis. Serum levels of high-density lipoprotein-cholesterol (HDL-C) and fasting blood glucose (FBG) were determined using Pars Azmun kits (Karaj, Iran) and the BT-3000 Auto-analyzer (Biotechnica, Rome, Italy) [26].

Serum levels of 25OHD were measured using an electrochemiluminescence method (ECL, Roche, Basel, Switzerland). Serum levels of 25OHD were measured using an assay with an intra- and inter-assay precision CV of 5.7% and 9.9%, respectively.

2.2.2. Serum prooxidant-antioxidant balance (PAB) assay

The serum PAB assay was used as previously described [15–17]. In brief, standard solutions were prepared by using variable proportions (0 to 100%) of 500 μM hydrogen peroxide (30%) (Merck, Massachusetts,

USA), that were added to 3 mM uric acid (in 10 mM NaOH). Then, 60 mg of 3,3',5,5'-Tetramethylbenzidine powder (TMB; Fluka) was dissolved in 10 mL DMSO. The TMB cation was prepared by adding 1 mL of the TMB/DMSO solution to 50 mL of acetate buffer (0.05 M buffer; pH 4.5). In the next step, 175 μL of chloramine T (100 mM; Applichem: A4331, Darmstadt, Germany) solution was added to 50 mL of acetate buffer (0.05 M buffer; pH 4.5). The solution was mixed and incubated for 1 h at room temperature in a dark place and shaken gently several times at the end of incubation. In the next step enzyme solution of peroxidase (16.5 μL ; Applichem: 230 U/mg, A3791,0005, Darmstadt, Germany) was added to TMB cation solution (50 mL), then the solution was placed into 1 mL aliquots and stored at -20°C . The TMB solution was prepared by mixing 200 μL of TMB/DMSO with 10 mL of acetate buffer (0.05 M buffer; pH 5.8), mixed well and placed in 4°C . Then, 10 μL of each sample, blank (distilled water) or standard samples were added to well of plate on ice. Working solution was ready by adding TMB cation (1 mL) to TMB solution (10 mL), and mixed well for 6 min (on a shaker) at room temperature in a dark place and used immediately. 200 μL of working solution added into the wells and incubated for 12 min in dark place at 37°C . After the incubation, 50 μL of HCl (2 N) was added to all wells, then the absorption was read by an ELISA reader at a wavelength of 450 nm together with a reference wavelength of 620 or 570 nm. A standard curve was presented by absorption readings of the standard samples. The values of the serum PAB are presented in Hamidi-Koliakos (HK) units, which show percentage of hydrogen peroxide in the standard solution. The values of PAB in the subject samples were then assessed via the values acquired from the above standard curve.

2.2.3. Statistical analysis

SPSS statistical package for Windows was used for all calculations (SPSS 20 software, SPSS Inc., Chicago, IL, USA). Statistical significance for the data compared in the adolescent girls before and after vitamin D administration was evaluated using the Student's *t*-test. Descriptive statistics including mean, frequency and standard deviation (SD) were determined for all variables and were expressed as mean $\pm$ SD for normally distributed variables. Data received for independent variables was analyzed using Student's *t*-test. All the analyses were two-sided and statistical significance was set at $P < 0.05$.

3. Results

3.1. Clinical and biochemical features

A comparison of the clinical and biochemical variables in adolescent girls pre- and post-intervention is summarized in Table 1 and was previously published [26]. The results show that serum PAB levels were increased ($P < 0.001$), whereas serum concentrations of LDL-C ($P < 0.001$), total cholesterol ($P < 0.001$) and HDL-C ($P < 0.01$) were reduced after vitamin D supplementation (Table 1).

There were no statistically significant differences in the clinical and biochemical factors among adolescent girls according to 25OHD categories at baseline ($P > 0.05$ for all variables; data have not been shown). Whereas, there were statistically significant improvements in weight ($P < 0.001$), BMI ($P < 0.001$) and FBG ($P = 0.02$) in adolescent girls who had 50–74.9 nmol/L serum 25OHD levels compared to those with a value < 50 nmol/L after vitamin D supplementation (Table 2).

Tables 3 and 4 show the comparison of the clinical and biochemical variables in adolescent girls according to BMI and HDL-C categories before and after vitamin D supplementation. The results demonstrate that vitamin D supplementation cause an increase in the serum PAB levels in both groups of adolescents with BMI < 25 and ≥ 25 (kg/m²) (Table 3). Moreover, vitamin D supplementation has increased the PAB levels in both groups of adolescents with HDL-C < 50 and ≥ 50 (mg/dl) (Table 4).

Table 1

Comparison of the clinical and biochemical variables in adolescent girls pre- and post-intervention.

Variables (mean $\pm$ SD) (n = 464)	Pre-intervention	Post-intervention	P-value
Weight (kg)	53 $\pm$ 11	54 $\pm$ 11	0.0001
BMI (kg/m ²)	21 $\pm$ 4	21 $\pm$ 4	0.31
WC (cm)	71 $\pm$ 9	70 $\pm$ 9	0.0001
Serum PAB (Hamidi-Koliakos (HK) units)	65 (38–103)	79 (45–122)	0.0001
SBP (mmHg)	99 $\pm$ 13	98 $\pm$ 13	0.37
DBP (mmHg)	65 $\pm$ 11	63 $\pm$ 11	0.003
FBG (mg/dl)	88 $\pm$ 11	85 $\pm$ 10	0.0001
Serum HDL-C (mg/dl)	47 $\pm$ 9	46 $\pm$ 8	0.003
Serum LDL-C (mg/dl)	101 $\pm$ 25	93 $\pm$ 21	0.0001
Serum Total cholesterol (mg/ dl)	163 $\pm$ 28	155 $\pm$ 26	0.0001
Serum Triglyceride	84 $\pm$ 37	82 $\pm$ 30	0.11
Serum 25OHD (nmol/l)	11 $\pm$ 10	38 $\pm$ 17	0.0001
Serum Hs-CRP (mg/l)	1 (0.5–1.6)	1 (0.5–1.8)	0.46
Serum Calcium (mmol/l)	9 (9.2–9.8)	10 (9.4–10.1)	0.0001
Serum Phosphate (mg/dl)	4 (3.6–4.2)	4 (3.8–4.3)	0.0001

Values expressed as mean $\pm$ SD. Between groups comparisons were assessed by nonparametric test for all non-normally distributed data (Wilcoxon test). BMI – body mass index; WC – waist circumference; PAB – prooxidant-antioxidant balance; SBP – systolic blood pressure; DBP – systolic blood pressure; FBG – fasting blood glucose; HDL-C – high-density lipoprotein cholesterol; LDL-C – low-density lipoprotein cholesterol; 25OHD – 25-hydroxyvitamin D; Hs-CRP – high sensitive C-reactive protein.

Table 2

Comparison of the clinical and biochemical variables in adolescent girls according to 25OHD categories after the vitamin D supplementation.

Variables (mean $\pm$ SD)	Serum 25OHD < 50 nmol/L (n = 341)	Serum 25OH 50–74.9 nmol/L (n = 106)	P-value
Weight (kg)	55 $\pm$ 11	51 $\pm$ 10	0.004
BMI (kg/m ²)	22 $\pm$ 4	20 $\pm$ 35	0.004
WC (cm)	70 $\pm$ 9	68 $\pm$ 8	0.06
Serum PAB (Hamidi- Koliakos (HK) units)	77 (43–121)	82 (46–130)	0.55
SBP (mmHg)	98 $\pm$ 13	99 $\pm$ 13	0.94
DBP (mmHg)	63 $\pm$ 11	63 $\pm$ 10	0.53
FBG (mg/dl)	86 $\pm$ 11	83 $\pm$ 10	0.02
Serum HDL-C (mg/dl)	46 $\pm$ 9	46 $\pm$ 7	0.61
Serum LDL-C (mg/dl)	95 $\pm$ 22	93 $\pm$ 22	0.48
Serum Total cholesterol (mg/dl)	157 $\pm$ 27	153 $\pm$ 26	0.11
Serum Triglyceride	84 $\pm$ 32	79 $\pm$ 29	0.07
Serum 25OHD (nmol/l)	30 $\pm$ 12	60 $\pm$ 6	0.0001
Serum Hs-CRP (mg/l)	1 (0.5–1.8)	1 (0.5–1.7)	0.63
Serum Calcium (mmol/l)	10 (9.4–10.0)	10 (9.4–10.1)	0.09
Serum Phosphate (mg/dl)	4 (3.8–4.3)	4 (3.8–4.3)	0.89

Values expressed as mean $\pm$ SD. Between groups comparisons were assessed by nonparametric test for all non-normally distributed data (Wilcoxon test). BMI – body mass index; WC – waist circumference; PAB – prooxidant-antioxidant balance; SBP – systolic blood pressure; DBP – systolic blood pressure; FBG – fasting blood glucose; HDL-C – high-density lipoprotein cholesterol; LDL-C – low-density lipoprotein cholesterol; 25OHD – 25-hydroxyvitamin D; Hs-CRP – high sensitive C-reactive protein.

3.2. Correlation of PAB with the clinical and biological parameters

The correlations between serum PAB with some other cardiovascular disease (CVD) risk factors among the adolescent girls are summarized in Tables 5 and 6. The results obtained from the current study show that there were no statistically significant associations between the serum PAB and any other biochemical factor parameters ($P > 0.05$ for all variables).

4. Discussion

To the best of our knowledge, this is the first study examining the effects of high dose vitamin D supplementation on PAB levels among adolescent girls. The results showed that vitamin D supplementation for 9 weeks caused a statistically significant decrease in serum levels of LDL-C, total cholesterol and HDL-C, and an increase in serum PAB values. Our findings are at variance with previous reports [27–29]. These showed that there was a statistically significant correlation between serum HDL-C and 25OHD before the intervention ($P = 0.002$), while, changes in serum HDL-C were not affected by supplementation ($P = 0.05$). In a cross-sectional study of 17,411 individuals, there was a positive association between serum HDL-C and serum 25OHD after adjustment for confounding factors in adults (25–84 years old) but in this study there was no intervention with vitamin D supplements [28]. Nikooyeh et al., assessed serum 25OHD and HDL-C in the summer and winter, and found that both increased significantly in summer, and again did not have an intervention arm. Moreover, they evaluated these factors in six different cities of Iran but used a small sample ($n = 530$ in total) [29].

Previous studies have measured separately the total pro-oxidant and anti-oxidant capacities [8]. Whilst, the present study used the serum PAB assay that measures pro- and antioxidant content simultaneously.

There is little evidence for the antioxidant properties of 25OHD. However, it has been reported that 25OHD can induce genes that can regulate redox balance, such as glutathione peroxidase, glucose-6-phosphate dehydrogenase and thioredoxin reductase [30,31].

It is reported that vitamin D has anticancer effects, perhaps via the effects of 25OHD binding to the vitamin D receptors that then translate to the nucleus where it binds to gene regulator regions [32–34]. However, the studies on the effects of 25OHD on the oxidative stress are limited. One study has shown that the prooxidative effect of 25OHD can reduce cancer development [18]. Whilst another study has reported that 25OHD can exert its anticancer effects through its antioxidative properties. A further study suggests that 25OHD may prevent prostate cancer through its transcriptional activation of glucose-6-phosphate dehydrogenase activity [35].

The results of the current study show that high dose vitamin D supplementation for a period of 9 weeks increased the serum PAB levels in adolescent girls. This result may be due to the high dose, or long-term of vitamin D supplementation in this study. There have been no other similar studies that have done a similar research among adolescents, but previous studies have documented that the regulatory mechanisms of 25OHD can be influenced by the treatment time, dosage, cell type, environment and tissue [18].

There were no statistically significant differences in the clinical and biochemical factors among adolescent girls according to 25OHD categories at baseline. Moreover, there were no statistically significant differences in PAB and HDL-C levels among adolescent girls with serum levels of 25OHD < 50 and 50–74.9 nmol/L after the supplementation. Whereas, the vitamin D supplementation caused improvement in the weight, BMI and FBG in adolescent girls who had 50–74.9 nmol/L serum 25OHD levels compared to < 50 nmol/L ones.

Our results show no statistically significant association between serum PAB or other biochemical factors. To further clarify the effects of high dose supplementation of vitamin D in adolescents, it may be important to measure the components that are measured as part of PAB assay such as albumin or uric acid, to assess the importance of this finding on factors that are more explicitly linked with cardiovascular risk [15]. In addition, more randomized controlled trials using different doses and a placebo group are required to determine the effects of vitamin D supplementation on the oxidative stress.

4.1. Study limitations

In the present study, there was no control group based on advice of

Table 3

Comparison of the clinical and biochemical variables in adolescent girls according to BMI categories before and after vitamin D supplementation.

Variables (mean $\pm$ SD)	Pre-intervention		P-value	Post-intervention		P-value
	BMI < 25 (kg/m ²) n = 376	BMI $\geq$ 25 (kg/m ²) n = 67		BMI < 25 (kg/m ²) n = 362	BMI $\geq$ 25 (kg/m ²) n = 67	
Weight (kg)	51 $\pm$ 8	71 $\pm$ 9	0.0001	51 $\pm$ 8	72 $\pm$ 9	0.0001
BMI (kg/m ²)	20 $\pm$ 3	28 $\pm$ 3	0.0001	20 $\pm$ 3	28 $\pm$ 2	0.0001
WC (cm)	69 $\pm$ 7	84 $\pm$ 9	0.0001	67 $\pm$ 6	83 $\pm$ 9	0.0001
Serum PAB (Hamidi-Koliakos (HK) units)	61 (39–98)***	79 (44–116)*	0.04	76 (44–125)	89 (5–120)	0.24
SBP (mmHg)	98 $\pm$ 13	105 $\pm$ 13	0.0001	98 $\pm$ 13	103 $\pm$ 13	0.002
DBP (mmHg)	65 $\pm$ 12	69 $\pm$ 10	0.006	63 $\pm$ 11	65 $\pm$ 10	0.10
FBG (mg/dl)	87 $\pm$ 11	88 $\pm$ 11	0.73	85 $\pm$ 11	86 $\pm$ 11	0.60
Serum HDL-C (mg/dl)	47 $\pm$ 9	46 $\pm$ 10	0.42	46 $\pm$ 8	46 $\pm$ 12	0.52
Serum 25OHD (nmol/l)	11 $\pm$ 10	9 $\pm$ 8	0.16	38 $\pm$ 17	36 $\pm$ 15	0.35
Serum Hs-CRP (mg/l)	1 (0.4–1.4)	2 (1.1–3.0)	0.0001	1 (0.4–1.5)	2 (1.0–2.7)	0.0001
Serum Calcium (mmol/l)	9 (9.2–9.8)	10 (9.2–9.8)	0.12	10 (9.4–10.1)	10 (9.4–10.1)	0.39
Serum Phosphate (mg/dl)	4 (3.7–4.2)	4 (3.6–4.1)	0.06	4 (3.8–4.3)	4 (3.8–4.3)	0.64

Values expressed as mean $\pm$ SD. Between groups comparisons were assessed by nonparametric test for all non-normally distributed data (Wilcoxon test). BMI - body mass index; WC - waist circumference; PAB - prooxidant-antioxidant balance; SBP - systolic blood pressure; DBP - systolic blood pressure; FBG - fasting blood glucose; HDL-C - high-density lipoprotein cholesterol; 25OHD - 25-hydroxyvitamin D; Hs-CRP - high sensitive C-reactive protein.

*** (P = 0.0001) and * (P = 0.013) are related to paired tests comparisons (Wilcoxon test) within BMI groups.

our ethics committee. Also, the high dose and short intervention period may have affected the results in this study compared with previous studies on vitamin D supplementation. Moreover, differences in serum 25OHD concentrations in different seasons were not considered. Although it seems that this issue cannot really affect the interpretation of our data, because the study was done over 9 weeks.

5. Conclusions

The results of the present study show that high dose vitamin D supplementation for a period of 9 weeks increases the serum PAB in adolescent girls. More randomized controlled trials with a control group warranted to assess the effects of vitamin D supplementation on the oxidative stress.

Conflict of interests

The authors declare no conflict of interests.

Financial disclosure

This study was supported by Mashhad University of Medical Sciences (MUMS), Iran (ID number 931188).

The author contribution

Study Design: Majid Ghayour-Mobarhan, Ameneh Timar, Maryam Saberi-Karimian

Data Collection: Ameneh Timar, Seyed Mohammad Reza Parizadeh, Reihaneh Sabbaghzadeh, Maryam Emadzadeh, Fatemeh Eshaghi

Statistical Analysis: Maryam Saberi-Karimian, Hamideh Ghazizadeh

Data Interpretation: Ameneh Timar, Maryam Saberi-Karimian, Majid Ghayour-Mobarhan, Gordon A. Ferns

Manuscript Preparation: Ameneh Timar, Maryam Saberi-Karimian, Majid Ghayour-Mobarhan

Literature Search: Ameneh Timar, Maryam Saberi-Karimian, Majid Ghayour-Mobarhan

Funds Collection: Majid Ghayour-Mobarhan

Table 4

Comparison of the clinical and biochemical variables in adolescent girls according to HDL-C categories before and after the vitamin D supplementation.

Variables (mean $\pm$ SD)	Pre-intervention		P-value	Post-intervention		P-value
	Serum HDL-C < 50 mg/dl (n = 262)	Serum HDL-C $\geq$ 50 mg/dl (n = 155)		Serum HDL-C < 50 mg/dl (n = 287)	Serum HDL-C $\geq$ 50 mg/dl (n = 111)	
Weight (kg)	54 $\pm$ 11	52 $\pm$ 11	0.02	54 $\pm$ 11	53 $\pm$ 10	0.31
BMI (kg/m ²)	22 $\pm$ 4	21 $\pm$ 3.84	0.01	21 $\pm$ 4	21 $\pm$ 4	0.64
WC (cm)	72 $\pm$ 9	70 $\pm$ 9	0.04	70 $\pm$ 9	69 $\pm$ 9	0.26
Serum PAB (Hamidi-Koliakos (HK) units)	65 (36.5–105)***	62 (38.1–90.5)***	0.63	80 (45.8–125.9)	76 (41.2–118.0)	0.34
SBP (mmHg)	99 $\pm$ 13	100 $\pm$ 12	0.40	98 $\pm$ 13	100 $\pm$ 13	0.25
DBP (mmHg)	65 $\pm$ 11	67 $\pm$ 11	0.35	63 $\pm$ 10	64 $\pm$ 12	0.70
FBG (mg/dl)	86 $\pm$ 12	89 $\pm$ 11	0.009	84 $\pm$ 10	88 $\pm$ 11	0.002
Serum 25OHD (nmol/l)	10 $\pm$ 9	12 $\pm$ 11	0.02	37 $\pm$ 17	39 $\pm$ 17	0.27
Serum Hs-CRP (mg/l)	1 (0.6–1.8)	1 (0.4–1.4)	0.08	1 (0.4–1.8)	1 (0.5–1.5)	0.93
Serum Calcium (mmol/l)	9 (9.2–9.8)	9 (9.3–9.8)	0.09	10 (9.3–10.0)	10 (9.4–10.3)	0.002
Serum Phosphate (mg/dl)	4 (3.6–4.2)	4 (3.7–4.2)	0.04	4 (3.8–4.3)	4 (3.9–4.4)	0.27

Values expressed as mean $\pm$ SD. Between groups comparisons were assessed by nonparametric test for all non-normally distributed data (Wilcoxon test). BMI - body mass index; WC - waist circumference; PAB - prooxidant-antioxidant balance; SBP - systolic blood pressure; DBP - systolic blood pressure; FBG - fasting blood glucose; 25OHD - 25-hydroxyvitamin D; Hs-CRP - high sensitive C-reactive protein.

*** (P = 0.0001) are related to paired tests comparisons within HDL-C groups.

Table 5
Correlation between PAB with some CVD risk factors among the adolescent girls before taking vitamin D₃ (n = 464).

	PAB arbitrary unit: HK	Serum TC (mg/ dl)	Serum TG (mg/ dl)	Serum HDL-C (mg/ dl)	Serum LDL-C (mg/dl)	Serum Hs-CRP (mg/l)	Serum Calcium (mmol/l)	Serum Phosphate (mg/ dl)
Serum PAB (Hamidi-Koliakos (HK) units)	1	0.03	0.02	-0.05	0.00	0.02	-0.02	0.00
	Pearson Correlation	0.60	0.61	0.31	0.99	0.75	0.68	0.98
	Sig. (2-tailed)	420	433	425	420	402	426	422
Serum Total cholesterol (mg/dl)	0.03	1	0.29**	0.36**	0.82**	0.00	0.22**	0.08*
	Pearson Correlation	0.60	0.0001	0.0001	0.0001	0.94	0.0001	0.02
	Sig. (2-tailed)	420	832	802	792	574	800	796
Serum Triglyceride (mg/dl)	0.02	0.29**	1	-0.09**	0.18**	0.15**	0.20**	0.16**
	Pearson Correlation	0.61	0.0001	0.007	0.0001	0.0001	0.0001	0.0001
	Sig. (2-tailed)	433	832	853	843	610	851	845
Serum HDL-C (mg/dl)	-0.05	0.36**	-0.09**	1	0.12**	-0.04	0.17**	0.08*
	Pearson Correlation	0.31	0.0001	0.007	0.001	0.36	0.0001	0.02
	Sig. (2-tailed)	425	853	853	841	591	826	828
Serum LDL-C (mg/dl)	0.00	0.82**	0.18**	0.12**	1	-0.03	0.22**	0.1**
	Pearson Correlation	0.99	0.0001	0.001	0.001	0.50	0.0001	0.002
	Sig. (2-tailed)	420	792	841	843	585	815	817
Serum Hs-CRP (mg/l)	0.02	0.00	0.15**	-0.04	-0.03	1	0.01	0.00
	Pearson Correlation	0.75	0.0001	0.36	0.50	0.88	0.93	0.93
	Sig. (2-tailed)	402	574	591	585	587	576	576
Serum Calcium (mmol/l)	-0.02	0.22**	0.20**	0.17**	0.22**	0.01	1	0.24**
	Pearson Correlation	0.68	0.0001	0.0001	0.0001	0.88	0.0001	0.0001
	Sig. (2-tailed)	426	851	826	815	587	844	844
Serum Phosphate (mg/dl)	0.00	0.08*	0.16**	0.08*	0.11**	0.00	0.24**	1
	Pearson Correlation	0.98	0.001	0.02	0.002	0.93	0.0001	0.0001
	Sig. (2-tailed)	422	845	828	817	576	844	851

PAB - prooxidant-antioxidant balance; HDL-C - high-density lipoprotein cholesterol; LDL-C - low-density lipoprotein cholesterol; Hs-CRP - high sensitive C-reactive protein.

** Correlation is significant at the 0.01 level (2-tailed).

* Correlation is significant at the 0.05 level (2-tailed).

Table 6
Correlation between the changes of PAB with some CVD risk factors among the adolescent girls at baseline and after 9 weeks of vitamin D supplementation, (n = 464).

	PAB arbitrary unit:		Serum TC(mg/dl)		Serum TG(mg/dl)		Serum HDL-C (mg/dl)		Serum LDL-C (mg/dl)		Serum Hs-CRP (mg/l)		Serum Calcium (mmol/l)		Serum Phosphate (mg/dl)	
	HK															
Serum PAB (Hamidi-Koliakos (HK) units)	1	Pearson Correlation Sig. (2-tailed)	–0.02	0.62	–0.02	0.64	–0.04	0.84	–0.01	0.46	–0.04	0.00	0.94	0.28	0.05	
Serum Total cholesterol (mg/dl)	463	N	389	1	401	385	378	0.73	0.38	367	0.00	394	388	0.05	0.28	
	0.62	Pearson Correlation Sig. (2-tailed)	–0.02	0.62	0.13	0.43	0.0001	0.0001	0.0001	0.91	0.0001	0.0001	0.0001	0.05	0.28	
Serum Triglyceride (mg/dl)	389	N	594	1	590	558	542	0.12	0.18	438	0.03	562	557	0.03	0.40	
	–0.02	Pearson Correlation Sig. (2-tailed)	0.13	0.001	1	0.03	0.51	0.005	0.005	0.55	0.18	0.0001	0.0001	0.40	0.07	
Serum HDL-C (mg/dl)	401	N	590	0.43	649	601	1	0.26	0.256	469	0.07	606	600	0.11	0.16	
	–0.04	Pearson Correlation Sig. (2-tailed)	0.43	0.0001	0.03	0.51	0.0001	0.0001	0.0001	–0.087	0.256	0.0001	0.0001	0.11	0.16	
Serum LDL-C (mg/dl)	385	N	558	0.73	601	605	605	0.26	0.37	445	0.02	585	588	0.06	0.16	
	–0.01	Pearson Correlation Sig. (2-tailed)	0.73	0.0001	0.12	0.26	0.0001	0.0001	0.0001	0.73	0.37	0.0001	0.0001	0.06	0.16	
Serum Hs-CRP (mg/l)	378	N	542	0.00	587	587	587	0.02	0.12	436	0.02	568	571	0.01	0.12	
	–0.04	Pearson Correlation Sig. (2-tailed)	0.00	0.91	–0.03	0.07	–0.09	0.02	0.02	1	0.12	–0.06	0.12	0.01	0.12	
Serum Calcium (mmol/l)	367	N	438	0.38	469	445	445	0.37	0.37	501	0.22	461	452	0.01	0.04	
	0.00	Pearson Correlation Sig. (2-tailed)	0.38	0.0001	0.18	0.26	0.0001	0.0001	0.0001	–0.06	0.22	1	0.33	0.01	–0.04	
Serum Phosphate (mg/dl)	394	N	562	0.05	606	585	585	0.06	0.12	461	0.06	635	624	0.04	0.33	
	0.05	Pearson Correlation Sig. (2-tailed)	0.05	0.28	0.03	0.07	0.11	0.06	0.16	461	0.12	–0.04	1	0.04	0.33	
	0.28	N	557	0.28	600	588	588	0.16	0.16	452	0.01	624	629	0.01	0.33	
	388															

PAB - prooxidant-antioxidant balance; HDL-C - high-density lipoprotein cholesterol; LDL-C - low-density lipoprotein cholesterol; Hs-CRP - high sensitive C-reactive protein.

** Correlation is significant at the 0.01 level (2-tailed).

* Correlation is significant at the 0.05 level (2-tailed).

Acknowledgements

The authors are thankful to the participants who contributed to this study.

References

- [1] McCarty DE. Resolution of hypersomnia following identification and treatment of vitamin D deficiency. Official publication of the American academy of sleep medicine. *JCSM* 2010;6(6):605.
- [2] Ma Y, Johnson CS, Trump DL. Mechanistic insights of vitamin D anticancer effects. *Vitam Horm* 2016(100):395–431.
- [3] Vanoirbeek E, Krishnan A, Eelen G, Verlinden L, Bouillon R, Feldman D, et al. The anti-cancer and anti-inflammatory actions of 1, 25 (OH) 2D3. *Best Pract Res Clin Endocrinol Metab* 2011;25(August (4)):593–604.
- [4] Wilmanski T, Zhou X, Zheng W, Shinde A, Donkin SS, Wendt M, et al. Inhibition of pyruvate carboxylase by 1 α , 25-dihydroxyvitamin D promotes oxidative stress in early breast cancer progression. *Cancer Lett* 2017;411(28):171–81.
- [5] Trump DL, Deeb K, Johnson CS. Vitamin D: considerations in the continued development as an agent for cancer prevention and therapy. *Cancer J* 2010;16(1):1.
- [6] Tangpricha V, Pearce EN, Chen TC, Holick MF. Vitamin D insufficiency among free-living healthy young adults. *Am J Med* 2002;112(8):659.
- [7] Lips P, Duong T, Oleksik A, Black D, Cummings S, Cox D, et al. A global study of vitamin D status and parathyroid function in postmenopausal women with osteoporosis: baseline data from the multiple outcomes of raloxifene evaluation clinical trial. *JCEM* 2001;86(3):1212–21.
- [8] Ginde AA, Liu MC, Camargo CA. Demographic differences and trends of vitamin D insufficiency in the US population, 1988–2004. *Arch Intern Med* 2009;169(6):626–32.
- [9] Langlois K, Greene-Finestone L, Little J, Hidioglou N, Whiting S. Vitamin D status of Canadians as measured in the 2007 to 2009 Canadian Health Measures Survey. *Health Rep* 2010;21(1):47.
- [10] Saki F, Dabbaghmanesh MH, Omrani GR, Bakhshayeshkaram M. Vitamin D deficiency and its associated risk factors in children and adolescents in southern Iran. *Public Health Nutr* 2015:1–6.
- [11] Ataie-Jafari A, Qorbani M, Heshmat R, Ardalan G, Motlagh ME, Asayesh H, et al. The association of vitamin D deficiency with psychiatric distress and violence behaviors in Iranian adolescents: the CASPIAN-III study. *J Diabetes Metab Disord* 2015;14(1):62.
- [12] Bonakdaran S, Fakhraee F, Karimian MS, Mirhafez SR, Rokni H, Mohebbati M, et al. Association between serum 25-hydroxyvitamin D concentrations and prevalence of metabolic syndrome. *ADVMS* 2016;61(2):219–23.
- [13] Halliwell B, Gutteridge J. Oxidative stress: adaptation, damage, repair and death. "Free radicals in biology and medicine". New York: Oxford University Press; 1999.
- [14] Habbous M, Herbeth B, Vincent-Viry M, Lamont JV, Fitzgerald PS, Visvikis S, et al. Serum total antioxidant status, erythrocyte superoxide dismutase and whole-blood glutathione peroxidase activities in the Stanislas cohort: influencing factors and reference intervals. *CCLM* 2003;41(2):209–15.
- [15] Alamdari DH, Paletas K, Pegiou T, Sarigianni M, Befani C, Koliakos G. A novel assay for the evaluation of the prooxidant–antioxidant balance, before and after antioxidant vitamin administration in type II diabetes patients. *Clin Biochem* 2007;40(3):248–54.
- [16] Hamidi Alamdari D, Ordoudi SA, Nenadis N, Tsimidou MZ, Koliakos G, Parizadeh SM, et al. Comparison of prooxidant-antioxidant balance method with crocin method for determination of total prooxidant-antioxidant capacity. *IJBMS* 2009;12(April (2)):93–9.
- [17] Alamdari DH, Ghayour-Mobarhan M, Tavallaie S, Parizadeh MR, Moohebbati M, Ghafoori F, et al. Prooxidant–antioxidant balance as a new risk factor in patients with angiographically defined coronary artery disease. *Clin Biochem* 2008;41(April (6)):375–80.
- [18] Gombart AF. Vitamin d: oxidative stress, immunity, and aging. CRC Press; 2012.
- [19] Ibi M, Sawada H, Nakanishi M, Kume T, Katsuki H, Kaneko S, et al. Protective effects of 1 α , 25-(OH) 2 D 3 against the neurotoxicity of glutamate and reactive oxygen species in mesencephalic culture. *Neuropharmacology* 2001;40(6):761–71.
- [20] Garçon E, Sindji L, Leblondel G, Brachet P, Darcy F. 1, 25-Dihydroxyvitamin D3 regulates the synthesis of γ -glutamyl transpeptidase and glutathione levels in rat primary astrocytes. *J Neurochem* 1999;73(2):859–66.
- [21] Wiseman H. Vitamin D is a membrane antioxidant Ability to inhibit iron-dependent lipid peroxidation in liposomes compared to cholesterol, ergosterol and tamoxifen and relevance to anticancer action. *FEBS Lett* 1993;326(1–3):285–8.
- [22] Koren R, Rocker D, Kotestiano O, Liberman UA, Ravid A. Synergistic anticancer activity of 1, 25-dihydroxyvitamin D 3 and immune cytokines: the involvement of reactive oxygen species. *J Steroid Biochem Mol Biol* 2000;73(3):105–12.
- [23] Rocker D, Ravid A, Liberman UA, Garach-Jehoshua O, Koren R. 1, 25-Dihydroxyvitamin D 3 potentiates the cytotoxic effect of TNF on human breast cancer cells. *Mol Cell Endocrinol* 1994;106(1):157–62.
- [24] Ravid A, Rocker D, Machlenkin A, Rotem C, Hochman A, Kessler-Ickson G, et al. 1, 25-Dihydroxyvitamin D3 enhances the susceptibility of breast cancer cells to doxorubicin-induced oxidative damage. *Cancer Res* 1999;59(4):862–7.
- [25] Sundaram S, Gewirtz D. The vitamin D3 analog EB 1089 enhances the response of human breast tumor cells to radiation. *Radiat Res* 1999;152(5):479–86.
- [26] Khayyatadeh SS, Mirmousavi SJ, Fazeli M, Abasalti Z, Avan A, Javandoost A, et al. High dose Vitamin D supplementation is associated with an improvement in several cardio-metabolic risk factors in adolescent girls: a nine- week follow up study. *Ann Clin Biochem* 2017(January (1)). <https://doi.org/10.1177/0004563217707784>. 4563217707784.
- [27] Grant WB, Boucher BJ. Genetic and non-genetic effects of increased sun and vitamin D exposure: role in the observed healthy changes in cardiometabolic risk factors in Iranian children. *Public Health Nutr* 2018(June):1–4.
- [28] Jorde R, Grimnes G. Exploring the association between serum 25-hydroxyvitamin D and serum lipids—more than confounding? *Eur J Clin Nutr* 2018;72(April (4)):526.
- [29] Nikooyeh B, Abdollahi Z, Hajifaraji M, Alavi-Majid H, Salehi F, Yarpasvar AH, et al. Healthy changes in some cardiometabolic risk factors accompany the higher summertime serum 25-hydroxyvitamin D concentrations in Iranian children: national food and nutrition surveillance. *Public Health Nutr* 2018(March):1–9.
- [30] Peehl DM, Shinghal R, Nonn L, Seto E, Krishnan AV, Brooks JD, et al. Molecular activity of 1,25-dihydroxyvitamin D3 in primary cultures of human prostatic epithelial cells revealed by cDNA microarray analysis. *J Steroid Biochem Mol Biol* 2004;92:131–41.
- [31] Lin R, Nagai Y, Sladek R, Bastien Y, Ho J, Petrecca K, et al. Expression profiling in squamous carcinoma cells reveals pleiotropic effects of vitamin D3 analog EB1089 signaling on cell proliferation, differentiation, and immune system regulation. *Mol Endocrinol* 2002;16:1243–56.
- [32] Ma Y, Johnson CS, Trump DL. Mechanistic insights of vitamin d anticancer effects. *Vitam Horm* 2016;100:395–431. <https://doi.org/10.1016/bs.vh.2015.11.003>.
- [33] Vanoirbeek E, Krishnan A, Eelen G, Verlinden L, Bouillon R, Feldman D, et al. The anti-cancer and anti-inflammatory actions of 1,25(OH)₂D₃. *Best Pract Res Clin Endocrinol Metab* 2011;25(August (4)):593–604. <https://doi.org/10.1016/j.beem.2011.05.001>.
- [34] Krishnan AV, Feldman D. Mechanisms of the anti-cancer and anti-inflammatory actions of vitamin D. *Annu Rev Pharmacol Toxicol* 2011;51:311–36. <https://doi.org/10.1146/annurev-pharmtox-010510-100611>.
- [35] Bao BY, Ting HJ, Hsu JW, Lee YF. Protective role of 1 α , 25-dihydroxyvitamin D3 against oxidative stress in nonmalignant human prostate epithelial cells. *Int J Cancer* 2008;122(12):2699–706.